

Summary Report: The Characteristics of Spanish Reservoirs
Data Files: CD (Sau physical and
Investigator: J. Armengol
Dept. Ecology. Fac. Biology. Univ. Barcelona
Avd. Diagonal, 645. 08028-Barcelona

Contract Number: N62558-02-M-6382
R&D Number: 9383EN07

Date: 11 April 2003

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 11 APR 2003		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE The Characteristics of Spanish Reservoirs				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Dept. Ecology. Fac. Biology. Univ. Barcelona Avd. Diagonal, 645. 08028-Barcelona				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

The Characteristics of Spanish Reservoirs

J. Armengol & R. Merçé
Dept. Ecology. Fac. Biology. Univ. Barcelona
Avd. Diagonal, 645. 08028-Barcelona

Sau Reservoir was first filled in 1963 in a middle stretch of the Ter River, as part of a multi-use scheme, including hydroelectric power, agricultural irrigation, domestic and industrial water supply to metropolitan areas, and also recreational activities. Since it was built, the reservoir has experienced a process of increasing eutrophication (Vidal and Om 1993), from moderately eutrophic during the first years after filling, to severe eutrophication in the late 1980s. Several human activities in the basin fed the process: intensive use of fertilizers, pig and stock farming development, proliferation of industrial areas and changes in land use (Sabater et al. 1990; Sabater et al. 1991; Vidal and Om 1993; Sabater et al. 1995; Espadaler et al. 1997). To improve the quality of water supplied by the reservoir, wastewater treatment plants were built in main urban and industrial areas during the early 1990's (Vidal and Om 1993).

Limnological characterization of Sau Reservoir started in 1963, conducted by the local water supply company ("Aigües Ter-Llobregat", ATLL). Since 1995, this study is executed in collaboration with the Department of Ecology at the University of Barcelona, under supervision of Dr. Joan Armengol. Thus, a good database is available on both changes in the water body and on incoming materials through main inflows (the Ter River and the Major River, accounting for 85% and 8% of land drained by Sau Reservoir, respectively). The basic sampling frequency is monthly. Besides in-situ measured descriptors (i.e. temperature, conductivity, turbidity, dissolved oxygen concentration, pH, redox potential, and water transparency), chemical analyses are conducted on samples taken at several depths. Analyses for nutrients include nitrate, nitrite, ammonia, total dissolved nitrogen, total nitrogen, reactive soluble phosphorus, total dissolved phosphorus, total phosphorus, and silicate. Other dissolved components are analyzed, i.e. carbonates and bicarbonates as alkalinity, and main anionic dissolved substances (chloride and sulphates). Scarce measures of sulphides and metals in deep water are also available. The planktonic community of the reservoir is sampled as phytoplankton in the 5 top meters and zooplankton in the whole column. Measures of chlorophyll are obtained both, as

pigments retained in a filter from water taken at different depths, and as profiles given by a field fluorimeter. Total suspended solids, ash-free suspended solids, and elemental carbon and nitrogen content of suspended solids are also analyzed.

Besides being the frame for the long-term limnological program, Sau Reservoir has been the field for a number of detailed limnological studies. Armengol and co-workers (Armengol et al. 1986; Armengol and Vidal 1988) gave early evidence of the increasing eutrophication process of the reservoir, focusing their studies in the phosphorus content of the sediments. Vidal and Om (1993) put data from the long-term study into an historical framework, reaching similar conclusions. Armengol et al. (1999) and Han et al. (2000) studied the reservoir as a longitudinal water reactor, both from the physical and chemical point of view. Finally, detailed information about the dynamics of the planktonic microbial community of the reservoir can be found in Sommaruga et al. (1995), Comerma et al. (2001), Simek et al. (1998; 2000; 2001), and Gasol et al. (2002).

As general remarks, Sau Reservoir is an 18.5 km long, 130 Hm³ capacity, canyon-shape eutrophic system. It receives nutrient loads that exceed in the present 10 g SRP·m⁻²·year⁻¹ and 500 g DIN·m⁻²·year⁻¹, supporting more than 250 mg chlorophyll-*a*·m⁻². Several hydrological patterns are described for the reservoir, from deep-water circulation during winter, to inter-flow during summer and autumn. The reservoir can be considered as a heterotrophic system in the riverine zone, where carbon imported from the river maintains a rich and high-productive bacterial community. The lacustrine zone, in contrast, develops a classical community maintained by the autochthonous production of phytoplankton.

Current research in the reservoir involves the study of the relevance of the meteorological forcing in the system, mainly focused on finding causal relationships to explain the short-term blooms of phytoplankton observed in the lacustrine area of Sau Reservoir. Also, a detailed historical research of the eutrophication process of the reservoir is being executed. This will include proposals for management targets at the watershed scale, applying both watershed and reservoir state-of-the-art models.

The study of Sau Reservoir limnology goes beyond the academic interest, as confirmed by the efforts from ATLL Water Supply Company to maintain the long-term work in the water body. Sau Reservoir is the main water supply of the city of Barcelona, and results generated during the long-term and detailed studies have been relevant to take management decisions that have improved the water quality supplied by the reservoir (i.e. construction of wastewater treatment plants in the basin, and correct management of reservoir outflows). Also, the social acceptance and use of the stored water is increasing. The hope is that this process will continue, as managers will implement correct measures to improve water quality, thanks to the results generated from limnological studies.

REFERENCES

- Armengol, J. and A. Vidal 1988. The use of different phosphorus fractions for the estimation of the trophic evolution of the Sau reservoir. *Arch. Hydrobiol. Beih.* 30: 61-70.
- Armengol, J., J.C. García, M. Comerma, M. Romero, J. Dolz, M. Roura, B.-H. Han, A. Vidal, and K. Simek 1999. Longitudinal Processes in Canyon Type Reservoirs: The Case of Sau (N.E. Spain), *In* [eds.], J.G. Tundisi and M. Straskraba *Theoretical Reservoir Ecology and its Applications*. International Institute of Ecology, Brazilian Academy of Sciences and Backhuys Publishers.
- Armengol, J., M. Crespo, J.A. Morguí, and A. Vidal 1986. Phosphorus budgets and forms of phosphorus in the Sau reservoir sediment: An interpretation of the limnological record. *Hydrobiologia* 143: 331-336.
- Comerma, M., J.C. García, J. Armengol, M. Romero, and K. Simek 2001. Planktonic food web structure along the Sau Reservoir (Spain) in Summer 1997. *International Review of Hydrobiology* 86: 195-209.
- Espadaler, I., J. Caixach, J. Om, F. Ventura, M. Cortina, F. Pauné, and J. Rivera 1997. Identification of organic pollutants in Ter river and its system of

reservoirs supplying water to Barcelona (Catalonia, Spain): A study by GC/MS and FAB/MS. *Wat.Res.* 31: 1996-2004.

Gasol, J.M., M.Comerma, J.C.García, J.Armengol, P.Kojecka, and K.Simek 2002. A trasplant experiment to identify the factors controlling bacterial abundance, activity, production and community composition in a eutrophic canyon-shaped reservoir. *Limnology and Oceanography* 47: 62-77.

Han, B., J.Armengol, J.C.García, M.Comerma, M.Roura, J.Dolz, and M.Straskraba 2000. The thermal structure of Sau reservoir (NE: Spain): a simulation approach. *Ecological Modelling* 125: 109-122.

Sabater, F., H.Guasch, E.Martí, J.Armengol, and S.Sabater 1995. The Ter: A Mediterranean river case-study in Spain, p. 419-438. *In* [eds.], C.E.Cushing, K.W.Cummins, and G.W.Minshall River and stream ecosystems. Elsevier.

Sabater, F., J.Armengol, and S.Sabater 1991. Physico-chemical disturbances associated with spatial and temporal variation in a Mediterranean river. *J.N.Am.Benthol.Soc.* 10: 2-13.

Sabater, F., S.Sabater, and J.Armengol 1990. Chemical characteristics of a mediterranean river as influenced by land uses in the watershed. *Wat.Res.* 24: 143-155.

Simek, K., J.Armengol, M.Comerma, J.C.García, P.Kojecka, J.Nedoma, and J.Hezlar 2001. Changes in the epilimnetic bacterial community composition, production, and protist-induced mortality along the longitudinal axis of a highly eutrophic reservoir. *Microbial Ecology* 42: 359-371.

Simek, K., J.Armengol, M.Comerma, J.C.García, T.H.Chrzanowski, M.Macek, J.Nedoma, and V.Straskraba 1998. Characteristics of protistan control of bacterial production in 3 reservoirs of different trophy. *International Review of Hydrobiology* 83: 485-494.

- Simek, K., K. Juergens, J. Nedoma, M. Comerma, and J. Armengol 2000.
Ecological role and bacterial grazing of *Halteria* spp.: small freshwater oligotrichs as dominant pelagic ciliate bacterivores. *Aquatic Microbial Ecology* 22: 43-56.
- Sommaruga, R., M. Krossbacher, W. Salvenmoser, J. Catalan, and R. Psenner 1995. Presence of large virus-like particles in a eutrophic reservoir. *Aquatic Microbial Ecology* 9: 305-308.
- Vidal, A. and J. Om 1993. The eutrophication process in Sau Reservoir (NE Spain): A long-term study. *Verh. Internat. Verein. Limnol.* 25: 1247-1256.